

Anaesthesia recommendations for **Dyke-Davidoff-Masson syndrome**

Disease name: Dyke-Davidoff-Masson syndrome

ICD 10: Q04.3

Synonyms: Cerebral hemi-atrophy; Cerebral hemi-hypoplasia

Disease summary: Dyke-Davidoff-Masson syndrome [1] is an extremely rare neurological condition with greater frequency in the paediatric population [3]. It was described by three clinicians in 1993, Dyke, Davidoff and Masson, who reported [9] patients with facial asymmetry, contralateral hemiparesis, seizures and mental retardation with pneumatoencephalographic findings on skull X-ray. There is a slight male [7] pre-ponderance. Adult cases [10] have also been reported, though very infrequently. The main causes of this rare syndrome are either congenital (in utero vascular occlusion) or acquired (perinatal hypoxia, intra-cranial haemorrhage and infections). Brain imaging [2] in the form of CT and MRI scans are required for diagnosis.

Literature regarding anaesthetic considerations for this syndrome is sparse.

Differential diagnoses [9] include Sturge-Weber syndrome, Fishman syndrome, Basal cell germinoma, Silver-Russell syndrome, Linear nevus syndrome and Rasmussen encephalitis.

Medicine is in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong



Find more information on the disease, its centres of reference and patient organisations on Orphanet: www.orpha.net

Typical surgery

Patients with Dyke-Davidoff-Masson syndrome can present for any type of elective or emergency surgery to the attending anaesthesiologist. Generally, these are paediatric patients [4]. In view of the occurrence of recurrent seizures and hemiparesis, they are prone to falls and hence develop orthopaedic injuries. Surgery may be required for fractures, non-unions, contractures or, sometimes, neurosurgical operations. Rarely, a few patients can undergo hemi-spherectomy for treatment-resistant seizures (success rate of 85% in carefully selected patients).

Type of anaesthesia

Anaesthesia for patients with Dyke-Davidoff-Masson syndrome can be quite challenging. There is not literature regarding anaesthetic considerations for this rare syndrome. In the presence of mental retardation, patient cooperation for regional anaesthesia and intravenous cannulation can be quite difficult to obtain. Both general and regional anaesthesia can be given with proper precautions. The preferred technique in such patients would be general anaesthesia, followed by ultrasound-guided regional nerve block.

In patients with developmental delays, venous cannulation can be challenging. The use of ultrasound-guidance for vascular access is recommended to improve success rates. Spinal deformities can also compound central neuraxial block performance and effect. Treatment medications taken for convulsions, hemiparesis and learning difficulties can have interactions with anaesthetic agents. Most anti-convulsants are enzyme inducers. Some patients may be taking drugs for psychiatric disorders like schizophrenia [8].

Necessary additional pre-operative testing (beside standard care)

If not already done before, neuroimaging [5] is a must for all these cases, which will show atrophy or hypoplasia of one of the cerebral hemispheres. History about brain insult in the foetal or early childhood period should be elicited, as well as maternal thrombophilia (in case of congenital cases) Other causes of mental retardation must be ruled out. An objective examination for the cause of seizures must be done and adequate epilepsy management must be started preoperatively. Besides standard pre-anaesthetic investigations, skull radiograph, contrast-computed tomographic (CECT) brain scan and electro-encephalography may be considered for evaluation. MRI (magnetic resonance imaging) of the brain can highlight changes in the cerebral hemisphere as well as bony structures, thus differentiating between acquired and congenital Dyke-Davidoff-Masson syndromes: the heterolateral brain structures are shifted toward the affected side in case of a congenital cause. Complete neurological assessment must be done to ascertain the level of hemiparesis (grade of motor/sensory loss). Serum anti-convulsant levels may be done in selected cases to ascertain adequate anti-epileptic medication level as well as to rule out their toxicity. Valproate can interfere with coagulation (fibrinogen) and a coagulation screening should ideally be performed before surgery at risk of bleeding. Special attention must be given to renal function tests and serum electrolytes.

This is part of the diagnostic setup of the syndrome: I suggest adding a section on diagnosis to include all these points. This should not be repeated before anaesthesia if the patient's medical file is OK.

Particular preparation for airway management

There is a possibility of difficult airway in view of facial asymmetry, ipsilateral hypertrophy of skull and sinus enlargement. In view of learning difficulties and mental retardation, awake intubation is not possible due to lack of patient cooperation. Inhalation anaesthesia or Intravenous sedation with preservation of spontaneous ventilation should be used for endotracheal intubation if a difficult airway is suspected or foreseen. Mask ventilation may be difficult due to improper or inadequate mask seal. Aspiration prophylaxis may be given in patients with recurrent or intractable seizures [6] and mental retardation. A difficult airway cart must be kept ready at all times and standard ASA guidelines must be followed for airway management.

Particular preparation for transfusion or administration of blood products

There are no known particular recommendations regarding transfusion or administration of blood products.

Particular preparation for anticoagulation

There is no known specific preparation for anticoagulation. Caution needs to be exercised in patients taking anticonvulsants and warfarin in view of enzyme induction.

Particular precautions for positioning, transportation and mobilisation

Peri-operative positioning needs special attention as these patients are mentally challenged and may have several facial/skull abnormalities. Documentation of pre-existing neuropathies must be done. All pressure points must be padded. Transportation and mobilisation must be gentle to prevent musculo-skeletal injuries.

Interactions of chronic disease and anaesthesia medications

There are no known disease-specific interactions with anaesthetic drugs. However, the medications taken by these patients can have interactions with anaesthetic agents, necessitating dose adjustments.

Anaesthetic procedure

Standard general anaesthesia followed by administration of regional nerve blocks (USG-guided) when appropriate is beneficial in these patients. Aspiration prophylaxis, preparation for difficult airway, securing a good venous access, ensuring adequate oxygenation and ventilation are the cornerstones of successful outcome. These children can have exaggerated separation anxiety and may have agitated behaviour [11] pre-operatively, leading to difficulty in induction. Extubation must also be cautious and expedited only after ensuring wakefulness and adequate tidal spontaneous ventilation. Post-operative care must be in a high-dependency care unit with requisite monitoring and involvement of the care-givers.

Particular or additional monitoring

Apart from standard ASA monitors, peripheral neuromuscular monitoring, and depth of anaesthesia monitor are recommended to prevent delayed reversal or awakening from anaesthesia. Due to the underlying cerebral anomalies and increased risk of convulsions, adequate depth of anaesthesia monitoring includes measuring its awake values (which can be challenging in this population) and interpreting any brisk increase as a possible sign of seizures. Invasive haemodynamic monitoring may be indicated in patients with other associated cardio-pulmonary abnormalities. Temperature monitoring must be meticulous to prevent the development of hypothermia in these vulnerable patients.

Possible complications

Complications can be disease-, anaesthesia- or procedure-related or due to surgical reasons. Special attention must be given to prevent hypoxia or hypercarbia, as these can trigger seizures. Some patients may develop emergence delirium or agitation and require prolonged hospital stay. The incidence of PONV (post-operative nausea vomiting) can be theoretically high, though no randomised studies have been done in this population. Pharmacologic PONV prophylaxis must be given in all patients. These patients must be dealt with compassion in view of their learning difficulties.

Post-operative care

These patients must be monitored in a high-dependency unit in the post-operative period. Vigilant observation must be done for seizures and anti-convulsant medications must be continued. Supplemental oxygen, adequate analgesia, psychological support and prevention of hypothermia must be ensured.

Disease-related acute problems and effect on anaesthesia and recovery

There are no specific interactions between disease-related acute problems and effect on anaesthesia and recovery. There could be delayed recovery from general anaesthesia due to the prolonged effects of anaesthetic agents, electrolytic disturbances or a post critical state following seizures under anaesthesia. It is necessary to use neuromuscular monitoring to ensure adequate reversal from muscle relaxants.

Ambulatory anaesthesia

Ambulatory anaesthesia is only advisable for short non-painful procedures. Standard precautions of postoperative care and peri-operative monitoring is advised. The presence of an expert care giver must be ensured before discharge. It is preferable to avoid day-care surgeries in view of seizure disorders, learning difficulties and maxillo-facial deformities.

Obstetrical anaesthesia

Literature is sparse regarding obstetric anaesthesia in patients with DDMS. Aspiration prophylaxis, adequate oxygenation, avoidance of supine-hypotension syndrome, continuation of anti-convulsants, and difficult airway preparedness must be achieved. It must be remembered that administering regional anaesthesia (central neuraxial block) may be extremely difficult in these patients due to lack of co-operation. Proper antenatal counselling must be done to develop a rapport with the patient to seek her co-operation for regional block, as it is much safer than general anaesthesia in an obstetric patient.

References

1. Yerdelen D, Zafer F. Dyke–Davidoff–Masson syndrome. *Neurosurg Quarterly* 2009;19:59–61
2. Singh P, Sagar K, Ahluwalia A. Dyke-Davidoff-Masson syndrome: Classical imaging findings. *J Pediatr Neurosci* 2010;5:124–125
3. Narain NP, Kumar R, Narain B. Dyke-Davidoff-Masson syndrome. *Indian Pediatr* 2008;45: 927–928
4. Thakkar PA, Dave RH. Dyke-Davidoff-Mason Syndrome: A rare cause of cerebral hemiatrophy in children. *J Pediatr Neurosci* 2016;11:252–254
5. Behera MR, Patnaik S, Mohanty AK. Dyke-Davidoff-Mason syndrome. *J Neurosci Rural Pract* 2012;3:411–413
6. Hsin YL, Chuang MF, Shen TW, Harnod T. Temporo-spatial analysis define epileptogenic and functional zone in a case of DDMS. *Seizure* 2011;20:713–716
7. Unal O, Tombul T, Cirak B, Anlar O, Incesu L, Kayan M. Left hemisphere and male sex dominance of cerebral hemiatrophy(DDMS). *Clin Imaging* 2004;28:163–165
8. Amann B, Garcia de la Iglesia C, Mckenna P, Pomarol-Clotel E, Sanchez-Guerra M, Orth M. Treatment- refractory Schizoaffective disorder in a patient with Dyke-Davidoff Masson Syndrome. *CNS Spectr* 2009;14:36–39
9. Koshy B, Surendrababu NR. Dyke-Davidoff-Masson Syndrome. *Ann Acad Med Singapore* 2010;39:501–502
10. Roy U, Panwar A, Mukherjee A, Biswas D. Adult presentation of Dyke-Davidoff-Mason Syndrome : A case report. *Case Rep Neurol* 2016;8:20–26
11. Lee JH, Lee ZI, Kim HK, Kwon SH: A case of Dyke-Davidoff-Masson syndrome in Korea. *Korean J Pediatr* 2006;49:208–211
12. Roy K, Talukdar A, Ray A, Pal P. Dyke-Davidoff-Masson syndrome-like picture in a case of Takayasu arteritis: an enigma. *BMJ Case Reports* 2012; Sep 11. DOI 10.1136/bcr-2012-006669
13. El-Bahri Ben Mrad F, Mrabet H, Ben Sghaier R, Mrabet A. Syndrome de Dyke-Davidoff-Masson: à propos de deux observations. *J Neuroradiol* 2005;32:50–53.

Date last modified: **April 2021**

This recommendation was prepared by:

Author

Uma Hariharan, Anaesthesiologist, Dr Ram Manohar Lohia Hospital and Post Graduate Institute of Medical Education & Research, CHS, New Delhi, India

Disclosure The author has no financial or other competing interest to disclose. This recommendation was unfunded.

This recommendation was reviewed by:

Reviewers

Francis Veyckemans, Anaesthesiologist, CHRU Lille, France
veyckemansf@gmail.com

Carlos R. Degrandi Oliveira, Anaesthesiologist, H Hospital Guilherme Álvaro, Santos, Brasil
degrandi@gmail.com

Please note that this recommendation has not been reviewed by an anaesthesiologist and a disease expert but by two anaesthesiologists instead.

Disclosure The reviewers have no financial or other competing interest to disclose.
