

Anaesthesia recommendations for **Lujan-Fryns syndrome**

Disease name: Lujan-Fryns syndrome

ICD 10: Q87.8

Synonyms: Lujan syndrome, X-linked mental retardation with marfanoid habitus, XLMR with marfanoid habitus

Disease summary: Lujan-Fryns syndrome (X-linked mental retardation with marfanoid habitus syndrome) is characterised by a marfanoid habitus becoming evident after puberty, typical craniofacial dysmorphism and behavioural problems. Recently mutations in the MED12 gene and in the UPF3B gene has already been described as the cause of Lujan-Fryns syndrome. Lujan-Fryns syndrome was first reported by Lujan in 1984 and Fryns offered other features of this syndrome in 1987. Craniofacial features in Lujan-Fryns syndrome include prominent forehead, long narrow face, small mandible, maxillary hypoplasia, long nose with high and narrow nasal bridge, short and deep philtra, thin upper lip, highly arched palate, receding chin, and low-set retroverted normal shaped ears. Complete or partial agenesis of corpus callosum, ascending aorta aneurysm and ventricular septal defects are other manifestations of Lujan-Fryns syndrome. The prevalence in the general population is not known but affects predominantly males. There is no available treatment for Lujan-Fryns syndrome, patients need special education and psychological support. Diaphragm hernia and pulmonary hypoplasia are the most common anomalies in this syndrome. Because of these anomalies. limited pulmonary reserves and marked facial dysmorphism make it difficult to have a secure airway. Furthermore, the presence of cardiovascular malformations can lead to significant problems during anaesthesia practice.

Medicine is in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong



Find more information on the disease, its centres of reference and patient organisations on Orphanet: www.orpha.net

Typical surgery

There is no typical surgery, but cleft palate surgery, vertebrae surgeries, cardiovascular surgeries can be possible due to accompanying anomalies.

Type of anaesthesia

There is no definite recommendation for either general or regional anaesthesia. However, considering the presence of craniofacial anomalies, which are specific features of this syndrome, the risk of difficult airway during general anaesthesia can make regional anaesthesia preferable.

Necessary additional pre-operative testing (beside standard care)

Most of clinical manifestations can be determined during physical examination but for the detection of cardiovascular anomalies such as ventricular septal defect and aortic aneurysm, echocardiogram should be part of the routine evaluation.

Particular preparation for airway management

Due to the presence of craniofacial anomalies, these patient are at high risk for a difficult airway.

Particular preparation for transfusion or administration of blood products

Not reported.

Particular preparation for anticoagulation

Not reported.

Particular precautions for positioning, transportation and mobilisation

Not reported.

Interactions of chronic disease and anaesthesia medications

Not reported.

Anaesthetic procedure

In these patients, no problems were encountered after using sevoflurane or propofol. However, adequate information about the use of other anaesthetic agents is not available.

Particular attention should be paid to the anaesthesia practice; it is the likely difficult airway that can be encountered in these patients. All necessary preparations must be made for the difficult airway.

Particular or additional monitoring

Standard monitoring of vital signs should be performed in all types of anaesthesia including sedation.

Possible complications

Complications due to difficult airway scenarios. Prepare for a difficult airway scenario to avoid a "Cannot intubate, cannot ventilate" situation.

Post-operative care

The degree of postoperative monitoring is depending on surgical procedure and preoperative condition of the patient.

Disease-related acute problems and effect on anaesthesia and recovery

As a last resort in a "Cannot intubate, cannot ventilate" situation, an emergency airway-puncture set must be at hand.

Ambulatory anaesthesia

Because of the possible difficult airway in patients with Lujan-Fryns syndrome, the needed additional devices to be prepared, general anaesthesia under ambulatory conditions or outside a major anaesthesiological department cannot be recommended.

Obstetrical anaesthesia

There are no obstetric case reports of patients with Lujan-Fryns syndrome.

References

1. Soni A, Meenawat A, Soni K, Srivastava V. Lujan Fryns Syndrome - A Case Report. *Clinical Practice* 2014;3(2):11–13
2. Basu S, Patil B, Aundhakar CD, Ingale SY. A rare case of Lujan Fryns syndrome. *International Journal of recent trends science technology* 2014; 9(3):341–342
3. Berry K, Mahajan S, Sahoo P, Cheeran S. Lujan Fryns Syndrome & Psychosis. *InternJ scientif study* 2014;2(5):105-107
4. Buggenhout GV, Fryns JP. Lujan Fryns Syndrome (mental retardation, X-linked, marfanoid habitus). *Orphanet J Rare Dis* 2006;1:26
5. De Hert M, Steemans D, Theys P, Fryns JP, Peuskens J. Lujan-Fryns syndrome in the differential diagnosis of schizophrenia. *Am J Med Genet* 1996;67:212–214
6. Dieckmann PM, Lucena LC, Dutra LA, Pedroso JL, Borsottini OGP. Marfanoid features and X-linked mental retardation associated with craniofacial abnormalities: lujan fryns syndrome. *Arq Neuropsiquiatr* 2013;71(1):66–69
7. Donders J, Toriello H, Van Doornik S. Preserved neurobehavioral abilities in Lujan-Fryns syndrome. *Am J Med Genet* 2002;107:243–246
8. Dotti MT, Malandrini A, Bartolini S, Fabrizi M, Federico A. Mental retardation with marfanoid syndrome: presentation of a family with different phenotypical expression. *Brain Dev* 1993;15:291–294
9. Santos ES, Aquino da Costa FK, Barros de Almeida DR, Ribeiro LG, Barbosa MM, Gomes RR, Angelim VML. Lujan-Fryns Syndrome – familial recurrence in the same generation. *Sci Med.* 2011;21(2):64–66
10. Fareed A. X-linked mental retardation with marfanoid habitus. *Catalogue for transmission genetics in Arabs* 2006;1–2
11. Fryns JP, Buttiens M, Van den Berghe H: Chromosome X-linked mental retardation and marfanoid syndrome. *J Genet Hum* 1988;36:123-128.
12. Fryns JP, Buttiens M. X-linked mental retardation with marfanoid habitus. *Am J Med Genet* 1987;28:267–274.
13. Fryns JP, Van den Berghe H. X-linked mental retardation with marfanoid habitus: a changing phenotype with age? *Genet Couns* 1991;2:241–244
14. Fryns JP. X-linked mental retardation with marfanoid habitus. *Am J Med Genet* 1991;38:233
15. Van Buggenhout G, Fryns JP. Lujan-Fryns syndrome (mental retardation, X-linked, marfanoid habitus). *Orphanet J Rare Dis* 2006;1:26
16. Gurrieri F, Neri G. A girl with the Lujan-Fryns syndrome. *Am J Med Genet* 1991;38:290–291
17. Khemka P, Basu M, Ray S, Mukhopadhyay S and Ghosh A. Lujan Fryns syndrome. *Sri Lanka journal of child health* 2012;41(4):201–202
18. Purandare KN, Markar TN. Psychiatric symptomatology of Lujan-Fryns syndrome: an X-linked syndrome displaying Marfanoid symptoms with autistic features, hyperactivity, shyness and schizophreniform symptoms. *Psychiatric Genetics*: 2005;15(3):229–231
19. Lacombe D, Bonneau D, Verloes A, Couet D, Koullischer L, Battin J: Lujan-Fryns syndrome (X-linked mental retardation with marfanoid habitus): Report of three cases and review. *Genet Couns* 1993;4:193–198
20. Lalatta F, Livini E, Selicorni A, Briscioli V, Vita A, Lugo F, Zollino M, Gurrieri F, Neri G. X-linked mental retardation with marfanoid habitus: First report of four Italian patients. *Am J Med Genet* 1991;38:228–232
21. Lerma-Carrillo I, Molina JD, Cuevas-Duran T, Julve-Correcher C, Espejo-Saavedra JM, Andrade-Rosa C, Lopez-Muñoz F. Psychopathology in the Lujan–Fryns syndrome: Report of two patients and review. *Am J Med Genet* 2006;140A:2807–2811
22. Lujan JE, Carlin ME, Lubs HA. A form of X-linked mental retardation with marfanoid habitus. *Am J Med Genet* 1984;17:311–322
23. Dündar NO, Dündar BN, Akkaya AE. A case of Lujan-Fryns syndrome. *Turk Arch Ped* 2010; 45: 291–294
24. Onal Ö, Dogru F, Aslanlar E, Celik JB. Anesthetic approach to the patient with Lujan-Fryns syndrome. 3rd International Conference on Surgery and Anesthesia, November 17-19, 2014 Chicago, USA
25. Dieckmann PM, Carneiro de Lucena L, Dutra LA, Pedroso JL, Orlando GP, Barsottini. Marfanoid features and X-linked mental retardation associated with craniofacial abnormalities: the Lujan-Fryns syndrome. *Arq Neuropsiquiatr* 2013;71(1):66–69

26. Pratap A, Agarwal A, Raja S, Khaniya S, Tiwari A, Kumar A. Fryns syndrome: A lethal mesoectodermal birth defect with variable expression in a pair of monozygotic twins. *Singapore Med J* 2007;48(4):e107
27. Khemka P, Basu M, Ray S, Mukhopadhyay S, Ghosh A. Lujan-Fryns Syndrome. *Sri Lanka Journal of Child Health* 2012;41(4):201–202
28. Purandare KN, Markar TN: Psychiatric symptomatology of Lujan-Fryns syndrome: an X-linked syndrome displaying Marfanoid symptoms with autistic features, hyperactivity, shyness and schizophreniform symptoms. *Psychiatr Genet* 2005;15:229–231
29. Rivera H, Ramírez-Dueñas MI, García-Ochoa C. Lujan syndrome in a Mexican boy. *Am J Med Genet* 1992;43:626–627
30. Basu S, Patil B, Aundhakar CD, Ingale SY. A Rare Case of Lujan Fryn Syndrome. *International Journal of Recent Trends in Science And Technology* 2014;9(3):341–342
31. Sargin M, Uluer MS, Sahin O. [Non-operating room anesthesia in a patient with Lujan-Fryns syndrome]. *Cukurova Med J.* 2016;41(1):33–35
32. Schwartz CE, Tarpey PS, Lubs HA, Verloes A, May MM, Risheg H, Friez MJ, et al. The original Lujan syndrome family has a novel missense mutation (p.N1007S) in the MED12 gene. *J Med Genet* 2007;44:472–477
33. Spaepen A, Hellemans H, Fryns JP. X-linked mental retardation with marfanoid habitus: the eye-catching psychiatric disorders [abstract]. *Am J Med Genet* 1994;51:611
34. Stathopulu E, Mackie Ogilvie C, Flinter FA. Terminal deletion of chromosome 5p in a patient with phenotypical features of Lujan-Fryns syndrome. *Am J Med Genet* 2003;119A:363–366
35. Stevenson RE, Schwartz CE, Schroer RJ: Lujan syndrome. In *XLinked Mental Retardation* Oxford, England; Oxford University Press; 2000:217–219
36. Van Buggenhout G; X-linked mental retardation with marfanoid habitus, Orphanet encyclopedia, September 2002. <https://www.orpha.net/data/patho/GB/uk-lujanfryns.pdf>
37. Wittine LM, Josephson KD, Williams MS. Aortic root dilatation in apparent Lujan-Fryns syndrome. *Am J Med Genet* 1999;86:405–409.

Date last modified: **August 2019**

This recommendation was prepared by:

Author(s)

Dr. Mehmet Sargin, Anaesthesiologist, Department of Anaesthesiology and Reanimation, Selcuk University Faculty of Medicine, Turkey
mehmet21sargin@yahoo.com

Disclosure The author has no financial or other competing interest to disclose. This recommendation was unfunded.

This recommendation was reviewed by:

Reviewer 1

Tino Münster, Anaesthesiologist, Department of anaesthesiology and intensive care medicine, Hospital Barmherzige Brüder, Regensburg, Germany
Tino.Muenster@barmherzige-regensburg.de

Reviewer 2

Özkan Onal, Anaesthesiologist, Associate Professor, Selcuk University Medical Faculty Department of Anaesthesiology and Intensive Care Medicine, Konya, Turkey
drozkanonal@gmail.com

Disclosures The reviewers have no financial or other competing interest to disclose.

Please note that this recommendation has not been reviewed by an anaesthesiologist and a disease expert but by two anaesthesiologists instead.
