

Anaesthesia recommendations for **Prader-Willi syndrome**

Disease name: Prader-Willi syndrome

ICD 10: Q87.1

Synonyms: Prader-Labhardt-Willi syndrome

Disease summary: Prader-Willi syndrome is a rare genetic disorder characterized by hypothalamic-pituitary abnormalities with severe hypotonia during the neonatal period and during the first two years of life, hyperphagia with a risk of morbid obesity during infancy and adulthood, learning difficulties and behavioural problems or severe psychiatric problems. The disease affects 1/25,000 births. The severe hypotonia at birth, which leads to sucking and swallowing problems and delayed psychomotor development, partially improves with age. Characteristic facial features (a narrow forehead, almond-shaped eyes, a thin upper lip and down-turned mouth), as well as very small hands and feet, are frequently observed. After this initial phase, the most striking signs appear including hyperphagia and absence of satiety often leading to severe obesity in affected children as young as two years of age. The situation may deteriorate quickly without adequate outside controls and obesity is a major factor influencing morbidity and mortality in these patients. Other associated endocrine abnormalities contribute to the clinical picture of short stature due to a growth hormone (GH) deficiency and incomplete pubertal development. Regularly, a decreased bone mineral density can be found without changed metabolism of calcium, phosphate, Vitamin D or parathyroid hormone. The degree of cognitive dysfunction varies widely from child to child. It is associated with learning disabilities, and impaired speech and language development that are aggravated further by psychological and behavioural troubles.

The disease is clinically and genetically heterogeneous. It is caused by anomalies involving the critical region of chromosome 15 (15q11-q13). The expert consensus is that diagnosis should be based on clinical criteria (Holm's criteria of 1993, revised in 2001) with confirmation by genetic analysis. Most cases are sporadic and familial recurrence is rare information that should be provided by genetic counselling. Management should be global and multidisciplinary. Early diagnosis, early multidisciplinary care and GH treatment have greatly improved the quality of life of affected children. There are currently no long-term data on the effect of GH treatment in adults, particularly concerning its effect on the behavioural problems and degree of autonomy obtained. In adults, complications linked to obesity and the issue of autonomy continue to pose important problems.

Medicine is in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong



Find more information on the disease, its centres of reference and patient organisations on Orphanet: www.orpha.net

Typical surgery

Orthopaedic surgery (e.g. spinal surgery for scoliosis); squint correction; cleft lip and palatal repair; dental treatment due to tooth decay; Orchidopexy (cryptorchism).

Type of anaesthesia

Both general and regional anaesthesia are challenging: using general anaesthesia may result in difficult airway management, landmarks for regional anaesthesia may be obscured due to morbid obesity.

Anaesthetic concerns of PWS include morbid obesity and sleep apnoea, difficult intravenous access, the potential for difficulties with airway management, risk for perioperative respiratory failure, primary myocardial involvement, aggressive and at times violent behaviour, convulsions, and disturbances in thermoregulation as well as glucose intolerance.

Necessary additional pre-operative testing (beside standard care)

Appropriate preoperative evaluation may depend on the presence of co-morbid conditions include in obesity-related complications – cardiovascular problems, diabetes mellitus, hypertension, sleep apnoea (**NOTE:** predisposition to cardiac-vascular complications independent of obesity).

- state of nutrition and hydration
- body mass index, neck circumference, dental issues
- salivary composition

Take care of a thorough evaluation of the patient's past and current respiratory status including Polysomnography as a part of an individualized preoperative workup in order to establish a baseline and identify those with severe OSA (as well as postoperative admission to the ICU for monitoring of respiratory status). The recordings include thoracic and abdominal breathing movements, nasal airflow, TC-PO₂, TC-CO₂, oxygen saturation, EEG, EOG (Electrooculography) and ECG.

Children with PWS frequently suffer from restrictive lung disease because of hypotonia, obesity and kyphoscoliosis. Preoperative evaluation with 12 lead ECG and echocardiography may be indicated in selected patients).

Particular preparation for airway management

In the perioperative management of a patient with Prader-Willi syndrome, special attention must be paid to the abnormalities in the upper and lower respiratory systems. Airway management may be complicated by the frequent association of poor dentition, micrognathia, palatal abnormalities, and limited neck mobility. The appropriate equipment to deal with the "cannot intubate/cannot ventilate" scenario should be available. Also, in patients in whom regional anaesthesia will be performed, there should be ready access to such equipment.

Intraoperative and postoperative respiratory compromise has also been a frequent perioperative problem in patients with PWS. These may include stridor, oxygen desaturation, hypercapnia, and intermittent bronchospasm with the need to use high peak inspiratory pressure.

Particular preparation for transfusion or administration of blood products

Not reported.

Particular preparation for anticoagulation

Not reported.

Particular precautions for positioning, transportation and mobilisation

Features that may have an impact on perioperative care include mental retardation with the propensity for aggressive behaviour.

Interactions of chronic disease and anaesthesia medications

There are no particular interactions, because there is no specific medication. Interactions depend on the presence of comorbid conditions and its drug therapy.

Anaesthetic procedure

Patients with PWS are presenting with muscular hypotension and therefore predisposed to gastric aspiration. The decreased motility of the gastrointestinal tract further heightens the risk of perioperative aspiration, and the sequel of aspiration may be more severe due to limited pulmonary reserves. The physiological set point of vomiting in PWS is abnormal, and these patients have a reduced tendency to vomit. Certainly the obese body habitus may be associated with a higher than normal incidence of hiatal hernia and increased intra-abdominal pressure. The perioperative aspiration risk is further exacerbated by a high incidence of rumination and lowered oesophageal sphincter tone. Rumination may contribute to the development of dental caries. Efforts should be made to reduce gastric acid secretion, increase intestinal motility, neutralize stomach contents, use body position to utilize gravity to reduce the tendency for passive regurgitation, secure the airway rapidly, decompress the stomach, and extubate the trachea cautiously. By these means, surgical morbidity and mortality related to aspiration of gastric contents may be reduced.

- nil per os for 6 hours
- peripheral intravenous cannula, topical anaesthetic cream should be used before ranitidine and metoclopramide intravenously, 1 hour prior to anaesthesia
- careful premedication with a sedative
- routine monitoring
- rapid sequence induction

Regional anaesthetic techniques play a role in such patients by eliminating the need for general anaesthesia and its perioperative risks. Regional anaesthesia can be helpful to provide intraoperative anaesthetic care and postoperative analgesia as a means of limiting the need for opioids and general anaesthetic agents. Both neuraxial techniques (epidural or spinal anaesthesia) and peripheral nerve blockade may be considered in these patients. Nevertheless, landmarks for regional anaesthesia may be obscured due to morbid obesity and the use of ultrasound is recommended to facilitate placement of the block and limit the incidence of complications.

A risk-benefit ratio has to be considered when using regional anaesthesia plus sedation in patients with PWS who may be at higher risk of aspiration than the general paediatric population. Sedation will likely be required not only for block placement, but also to ensure a cooperative patient during the surgical procedure.

Moreover, the syndrome may result in a prolonged and exaggerated response to every sedative and analgesic agent. It has been suggested that regional techniques and NSAIDs beneficial as a means of limiting opioid use in the immediate postoperative period.

The use of non-depolarizing neuromuscular blocking agents for muscle relaxation should be restricted because of potential for long-lasting neuromuscular blockade. Hypotonia has prompted some authors to caution against the use of neuromuscular blocking agents. However, several reports have demonstrated the safe use of various non-depolarizing neuromuscular blocking agents including pancuronium, atracurium, vecuronium, and rocuronium without evidence for prolonged effects. Neuromuscular monitoring is necessary and residual neuromuscular blockade should be promptly antagonized with neostigmine and glycopyrrolate. Although they would suggest caution with the use of succinylcholine in the presence of hypotonia given the theoretical risk of an exaggerated hyperkalaemic response, several of the reports have also demonstrated the safe use of succinylcholine in patients with PWS.

Some authors chose ketamine for its limited effects on respiratory function as well as its ability to provide both sedation and analgesia. Especially in older children, ketamine should be co-administered with either propofol or a benzodiazepine to limit the potential of emergence phenomena. Ketamine can be used with caution as its effects on the seizure threshold are controversial.

Propofol, isoflurane, sevoflurane and N₂O have been used for the maintenance of anaesthesia.

Other common problems include difficult venous puncture, food-seeking, behaviour, disturbances in thermoregulation, diabetes mellitus, arrhythmia, and cor pulmonale.

Particular or additional monitoring

Monitoring of the neuromuscular blockade is recommended.

Blood glucose and body temperature should be monitored carefully.

Additional monitoring depends on identification of elevated risk for intraoperative or postoperative problems due to comorbidity.

Possible complications

PWS patients tend to develop severe respiratory infections as a result of aspiration due to hypotonia leading to poor pharyngeal coordination, chronic aspiration and a weak cough.

Strategy for the prevention of perioperative complications for PWS patients:

- elective surgery should be postponed until complete recovery from a URI
- special attention must be paid to those patients complicated with severe OSA
- the increased risk of aspiration and postoperative apnoea should be considered.

Post-operative care

For close observation and therapy postoperative intensive care medicine is highly recommended to prevent sleep-related respiratory complications – even if neither narcotics nor intermediate or long lasting neuromuscular blocking agents are administered.

Postoperatively, emergence often is slow and accompanied by upper airway obstruction and snoring. Complications may include an increased number of apnoeic episodes, higher hypercapnic thresholds, obstructive episodes, bronchospasm and oxygen requirement.

Wound healing and mobilization can be slow.

Disease-related acute problems and effect on anaesthesia and recovery

The accompanying morbid obesity, OSA, pathologic central control of ventilation and excessive daytime sleepiness seems to be characteristic for PWS, and may be related to problems with the sleep-wake rhythm and hypothalamic dysfunction.

Patients with PWS present with Mental retardation with the propensity for aggressive behaviour, psychological disturbances, and obsessive behavioural problems.

In later childhood and adolescence, neurologic manifestations, endocrine signs and dysmetabolic abnormalities can occur.

Additional psychological disturbances, mental retardation and obsessive behavioural problems are frequent findings.

Cardiac conduction defects and primary myocardial involvement have been noted and convulsions are common.

Ambulatory anaesthesia

Ambulatory anaesthesia is not recommended.

Obstetrical anaesthesia

Not reported.

References

1. Dearlove OR, Dobson A, Super M. Anaesthesia and Prader-Willi syndrome. *Paediatr Anaesth* 1998;8:267–271
2. Burman P, Ritzén EM, Lindgren AC. Endocrine dysfunction in Prader-Willi syndrome: a review with special reference to GH. *Endocr Rev* 2001;22(6):787–99
3. Warwick JP, Mason DG. Obstructive sleep apnoea syndrome in children. *Anaesthesia* 1998;53:571–579
4. Curfs LM, Fryns JP. Prader-Willi syndrome: a review with special attention to the cognitive and behavioral profile. *Birth Defects Orig Artic Ser* 1992;28:99–104
5. Eiholzer U. Deaths in children with Prader-Willi syndrome. A contribution to the debate about the safety of growth hormone treatment in children with PWS. *Horm Res* 2005;63:33–39
6. Nixon GM, Brouillette RT. Sleep and breathing in Prader-Willi syndrome. *Pediatr Pulmonol* 2002;34:209–217
7. Legrand R, Tobias JD. Anesthesia and Prader-Willi syndrome: preliminary experience with regional anesthesia. *Paediatr Anaesth* 2006;16:712–722
8. Papavramidis ST, Kotidis EV, Gamvros O. Prader-Willi syndrome-associated obesity treated by biliopancreatic diversion with duodenal switch. Case report and literature review. *J Pediatr Surg* 2006;41:1153–1158
9. Camfferman D, McEvoy RD, O'Donoghue F, Lushington K. Prader Willi Syndrome and excessive daytime sleepiness. *Sleep Med Rev* 2008;12:65–75
10. Sloan TB, Kaye CI. Rumination risk of aspiration of gastric contents in the Prader-Willi syndrome. *Anesth Analg* 1991;73:492–495
11. Mantadakis E, Spanaki AM, Geromarkaki E, Vassilaki E, Briassoulis G. Near demise of a child with Prader-Willi syndrome during elective orchidopexy. *Paediatr Anaesth* 2006;16:790–793
12. Tseng CH, Chen C, Wong CH, Wong SY, Wong KM. Anesthesia for pediatric patients with Prader-Willi syndrome: report of two cases. *Chang Gung Med J* 2003;26:453–457
13. Rinaldi S, Rizzo L, Di Filippo A, Secchi S, Paternoster G, La Torre MS, Pascente C, Stanzani MR. Monopharmacologic general anaesthesia with sevoflurane in paediatric patient with Prader-Willi syndrome. *Minerva Anestesiol* 2002;68:783–790
14. Ostermeier AM, Hofmann-Kiefer K, Schwender D. Induction of anesthesia for a patient with sleep apnea syndrome *Anaesthesist* 2000;49:317–320
15. Silva PS, Monteiro Neto H, Andrade MM, Neves Cv. Negative-pressure pulmonary edema: A rare complication of upper airway obstruction in children. *Pediatr Emerg Care* 2005;21:751–754
16. Shine NP, Coates HL, Lannigan FJ, Duncan AW. Adenotonsillar surgery in morbidly obese children: Routine elective admission of all patients to the intensive care unit is unnecessary. *Anaesth Intensiv Care* 2006;36:724–730
17. Shapiro F, Sethna N. Blood loss in pediatric spine surgery. *Eur Spine J* 2004;13. Suppl 1:S6–17
18. Aravindan A, Singh AK, Kurup M, Gupta S. Anaesthetic management of paediatric patient with Prader-Willi syndrome for bariatric surgery. *Indian J Anaesth.* 2020;64:444–445
19. Tan HL, Urquhart DS. Respiratory Complications in Children with Prader Willi Syndrome. *Paediatric Respiratory Reviews* 22 (2017);52–5954
20. Crockett DJ, Ahmed SR, Sowder DR, Wootten CT, Chinnadurai S, Goudy SL. Velopharyngeal dysfunction in children with Prader-Willi syndrome after adenotonsillectomy. *Int J Pediatr Otorhinolaryngol* 2014;78:1731–1734. DOI: 10.1016/j.ijporl.2014.07.034
21. Pacilli M, Heloury Y, O'Brien M, Lioni T, Rowell M, Hutson J. Orchidopexy in children with Prader-Willi syndrome: Results of a long-term follow-up study. *J Pediatr Urology* 2018;14,1:63.e1-63.e6.

Date last modified: **August 2020**

This recommendation was prepared by:

Author

Heike Rakow, Anaesthesiologist, HELIOS Hospital, Schwerin, Germany
heike.rakow@helios-kliniken.de

Disclosure The author has no financial or other competing interest to disclose. This recommendation was unfunded.

This recommendation was reviewed by:

Reviewers

Joseph D. Tobias, Anaesthesiologist, Nationwide Children's Hospital, Columbus, USA
Joseph.Tobias@Nationwidechildrens.org

Bernd Schenk, Paediatrician (endocrinology and diabetology), HELIOS Hospital, Schwerin, Germany
bernd.schenk@helios-kliniken.de

Editorial Review 2020

Tino Münster, Anaesthesiologist, Department of anaesthesiology and intensive care medicine, Hospital Barmherzige Brüder, Regensburg, Germany
Tino.Muenster@barmherzige-regensburg.de

Disclosure The reviewers have no financial or other competing interest to disclose.
