

Anaesthesia recommendations for **Sotos syndrome**

Disease name: Sotos syndrome

ICD 10: Q87.3

Synonyms: Cerebral gigantism syndrome, Malan syndrome (Sotos syndrome 2)

Disease summary: Sotos syndrome is characterised by the presence of excessive growth during childhood, advanced bone age, macrocephaly, characteristic facial appearance and non-progressive learning difficulties [1]. It affects males and females in equal numbers. Its incidence is one in 14,000 live births. It was first described in 1964 by Dr. J.F. Sotos [2].

Until the early 2000s, diagnosis of Sotos syndrome was based upon the presence of characteristic clinical features as listed above [3]. It was subsequently discovered that mutations and deletions of the NSD1 (nuclear receptor-binding SET domain) gene – localised on chromosome 5q35.3, which codes for a histone methyltransferase implicated in transcriptional regulation – is responsible for more than 75 % of cases of Sotos syndrome [4]. In Europe and the USA, NSD1 mutations cause 60–80 % of cases of Sotos syndrome, whereby microdeletions of NSD1 cause approximately 10 % of cases. In contrast: in Japan, NSD1 microdeletions are the primary cause of Sotos syndrome in over 50 % of cases [4].

The function of the NSD1 gene has not yet been fully described. The gene codes for a histone methyltransferase, which acts as a transcriptional intermediary factor capable of both negatively and positively influencing transcription. NSD1 has been described as a “corepressor of genes that promote growth” [5]. While it remains unclear exactly how the malfunction of NSD1 leads to the features of Sotos syndrome, it is thought that mutations and microdeletions involving NSD1 disrupts the activity of genes involved in normal growth and development [4]. While over 95 % of cases are sporadic in aetiology, autosomal dominant inheritance has also been described in several cases [4].

As other genetic abnormalities were identified in patients with Sotos syndrome, patients with abnormalities of the NSD1 gene were later termed ‘Sotos syndrome 1’. Heterozygous mutations in the NFIX (Nuclear Factor I, X type) on chromosome 19p13.3 were identified in a cohort of children with the Sotos syndrome phenotype. They were labelled ‘Sotos syndrome 2’ [6]. Furthermore, in 2015, a loss-of-function, frameshift mutation in the APC2 (Adenomatous Polyposis Coli 2) gene was identified in two siblings with features of Sotos syndrome but without NSD1 mutations [7]. They exhibited intellectual disability, abnormal brain structure and typical facial features but no other features such as bone or heart abnormalities. The APC2 gene is a downstream regulator of the NSD1 gene and its expression is downregulated by abnormalities of the NSD1 gene, potentially explaining the resulting neurological manifestations [7]. This genotype-phenotype combination is now known as ‘Sotos syndrome 3’.

In Sotos syndrome, the childhood growth is particularly advanced in the first year of life, after which it stabilises, before normalising in puberty. Height measurements are consistently above the 97th percentile in years 2–6, while final height is usually in the high normal range [4].

Craniofacial features are distinctive. The forehead tends to be overly prominent in infancy, while in adolescences an elongated face with a prominent chin is seen. Hypertelorism and down slanting palpebral fissures are common, as are receding hairline, prominent jaw, high arched palate, anteverted nostrils and long ears. Occipitofrontal circumference remains between the 98th and 99.6th percentile throughout life [8].

Most patients have a non-progressive neurological dysfunction, though the degree of learning disability appears to be extremely variable. Delay in motor development and expressive language is common [4].

Other manifestations of Sotos syndrome are variable. Delayed attainment of milestones of development is to be expected. Developmental delay is present in 80–85 % of the patients, for example excessive drooling; central nervous system abnormalities: enlarged ventricles, increased subarachnoid spaces that may require treatment, agenesis or hypoplasia of the corpus callosum, agenesis of the septum pellucidum, hypoplasia and atrophy of the cerebellar vermis, large cisterna magna and abnormalities of the Sylvian fissure. Among musculoskeletal abnormalities, hypotonia is the most frequent (84 %), pectus excavatum or carinatum has also been reported. Ophthalmologic manifestations are frequent: strabismus, nystagmus, retinal and optic nerve anomalies [9]. Hypothyroidism, hyperthyroidism, hypoparathyroidism and thyrotoxicosis has been reported [9,10]. There is an increased incidence of different tumours at a young age [9,11], non-progressive hypotonia [12]. About 14 % of patients present with auditive-related conditions [10] such as hearing loss, EMO, recurrent ear infections, cholesteatoma, and degenerative changes of the eardrum. About 24% of individuals have congenital malformations or abnormalities of the head and neck [10], including high arched palate, auricular dysplasia, macroglossia, cleft lip and palate, and alveolar cleft. Feeding difficulties have also been reported [10]. It can be due to the hypotonia that may result in weakness in the muscles of swallowing and subsequent feeding difficulties or the associated congenital malformations of the head and neck. About 16 % of individuals were reported to have other conditions, including speech disorder, respiratory difficulties, laryngomalacia, severe OSA, gastroesophageal reflux disease, and parotitis [109].

Scoliosis and kyphoscoliosis are present in approximately 30 % of patients. Early detection of scoliosis is important to facilitate early intervention with back braces and/or surgery [8]. There is no reported or published data on pulmonary function capacity with the subset of patients with scoliosis compared to Sotos syndrome patients without scoliosis. Congenitally dislocated hips, genu vara/valga and propensity to fractures with minimal trauma have been described [8]. Childhood seizures are common in 50 %, frequently febrile in nature [4]. Behaviour abnormalities include social inhibition and attention deficit [13]. An increased occurrence of congenital heart disease is seen, the most common defects being patent ductus arteriosus and septal defects [8]. Renal anomalies (bifid, duplex, cystic or absent kidneys; vesicoureteral reflux; pelvoureteric junction obstruction) and genital anomalies (hypospadias; cryptorchidism) are present in 15 % of children with NSD1 abnormalities [14]. Recurrent upper respiration tract infections and otitis media are also common, and constipation also often requires treatment [4].

Medicine is in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong



Find more information on the disease, its centres of reference and patient organisations on Orphanet: www.orpha.net

Typical surgery

Cardiac, neuro, ENT and urogenital surgery.

Nissen fundoplication with/without gastrostomy.

Type of anaesthesia

There are no definite recommendations regarding either general anaesthesia or regional anaesthesia for patients with Sotos syndrome.

The vast majority of published cases describes the use of general anaesthesia for procedures for Sotos syndrome patients. Regional anaesthesia is reported in Sotos syndrome. Neurostimulator and ultrasound guidance should be used to reduce the risk of complications. Most of the children with Sotos syndrome are uncooperative and possibly aggressive, so regional anaesthesia should be performed after GA induction [11]. One case report of regional anaesthesia has been published in a patient diagnosed with Sotos syndrome and sustaining mild hypotonia. No complications were observed [15].

Necessary additional pre-operative testing (beside standard care)

A detailed pre-operative assessment and an understanding of the comorbidities of patients with Sotos syndrome is essential for the anaesthetic management. Patients with Sotos syndrome exhibit a high incidence of heart defects and genitourinary abnormalities, so detailed assessment for cardiac and renal disease should be performed pre-operatively.

Echocardiogram should be performed to assess for congenital heart defects, the most common being septal defects and patent ductus arteriosus. Renal ultrasound should also be considered if renal function is decreased.

Haematological investigations should include full blood count, urea, creatinine, electrolytes and coagulation profile.

Kyphoscoliosis may cause restrictive lung disease and require pre-operative respiratory testing [11].

History of seizures should be documented, together with the anticonvulsive drugs and the possible interactions with the anaesthetic drugs.

Anaesthetic procedure

- Consider the option of an inhalation induction and intravenous catheter placement when the patient is already unconscious.
- Oral sedative premedication can be considered but should be individualised depending on patient conditions such as hypotonia, OSA, or respiratory difficulties.
- Drugs decreasing the seizure threshold should be avoided, even if there is no history of seizure disorder; many children with Sotos syndrome have EEG abnormalities without clinical seizures [11].

- For general anaesthesia, parental presence during induction has been described as invaluable in achieving sufficient patient cooperation [16]. Tolerance of needles is variable, so oral sedative premedication can be considered [17].
- Despite the presence of craniofacial abnormalities including high arched palate, pointed chin and abnormally erupted teeth, endotracheal intubation has been performed without difficulty [16,18,19]. There are no documented cases of difficult intubation in the literature to date.
- Concern has been expressed regarding the use of muscle relaxants in patients with Sotos syndrome due to hypotonia. Muscle relaxants have been used in cases of Sotos syndrome without complications: succinylcholine, atracurium and vecuronium. It has been surmised that patients with Sotos syndrome do not have any significant aberrant clinical pharmacologic effects with the use of muscle relaxants [16,17,18]. Furthermore, the recovery trend of TOF post muscle relaxant administration appears to have been normal in the data to date [16].

Particular preparation for airway management

As detailed above, despite airway management concerns due to the craniofacial abnormalities associated with Sotos syndrome, the intubation in each case detailed in the literature to date has been straightforward [16,18,19]. Care should be taken, however, all the more so as a comprehensive airway assessment including Mallampati score is often not possible due to poor co-operation [16].

Of noted it has been reported that a depth of insertion of the orotracheal tube needs to be 3 cm deeper than that obtained through the standard formula ($12 + \text{age}/2$).

A previous case report had also demonstrated deeper endotracheal tube insertion in a child with Sotos syndrome. However, the tracheal tube sizes are appropriate for age, as predicted by usual formulae [11,12].

Macroglossia has been described [10].

Particular preparation for transfusion or administration of blood products

No particular concerns have been described.

Particular preparation for anticoagulation

No particular concerns have been described, but drug dosing should be adjusted if significant renal impairment is present.

Particular precautions for positioning, transportation and mobilisation

Extra care should be taken in positioning due to the above average height of these patients and laxity of joints. Head positioning may be difficult due to large occiput size [3]. These patients are prone to have fractures with minimal trauma.

Interactions of chronic disease and anaesthesia medications

Anaesthetic agents which depend on renal excretion should be used with caution if renal impairment is present.

It may be better to choose anaesthetic agents that increase the seizure threshold due to the high prevalence of seizures in Sotos syndrome [16].

Seizures in Soto syndrome should be managed with standard protocols for anti-seizure medication. In drug resistant seizures, ketogenic diet and vagal nerve stimulator could also be considered [16].

Particular or additional monitoring

No definitive guidelines are available. The presence of cardiovascular, renal and cerebrovascular disease often associated with Sotos syndrome should guide the need for invasive monitoring with arterial and central venous pressures on a case by case basis.

Possible complications

None of note documented to date.

Post-operative care

Post-operative care should depend on the type of surgery and the clinical situation of the patient post-operatively. Presence of a parent/guardian during the post-operative period has been shown to greatly increase cooperation of patients [16].

Monitor the appearance of possible episodes of respiratory failure. Severe OSA has been described [10].

Disease-related acute problems and effect on anaesthesia and recovery

No acute or emergency situations have arisen from the documented cases of anaesthesia for patients with Sotos syndrome to date.

Ambulatory anaesthesia

Day case anaesthesia has not been described. Due to likelihood of distress caused by unfamiliar surroundings during an inpatient stay, it is reasonable to consider ambulatory anaesthesia when possible for small procedures.

Obstetrical anaesthesia

There are cases of patients with Sotos syndrome giving birth, but no cases describing their obstetrical anaesthesia management have been described to date. There are no known case reports of the use of regional anaesthesia in Sotos syndrome. There is no known contra-

indication; however, the presence of scoliosis and cognitive impairment could make consent and performing neuroaxial anaesthesia difficult.

References

1. Cole TR, Hughes HE. Sotos syndrome: a study of the diagnostic criteria and natural history. *J Med Genet* 1994;31:20–32. DOI:10.1136/jmg.31.1.20
2. Sotos JF, Dodge PR, Muirhead D, Crawford JD, Talbot NB. Cerebral Gigantism in Childhood. A Syndrome of Excessively Rapid Growth and Acromegalic Features and a Nonprogressive Neurologic Disorder. *N Engl J Med* 1964;271:109–116. DOI:10.1056/NEJM196407162710301
3. Chierichini A, Messina A, Vergari A, Santoprete S, Rasanito L. Regional anesthesia in a child with sotos syndrome. *Int J Immunopathol Pharmacol* 2011;24:21–23. DOI:10.1177/03946320110241S205
4. Baujat G, Cormier-Daire V. Sotos syndrome. *Orphanet J Rare Dis*. 2007;2:36. DOI:10.1186/1750-1172-2-36
5. Türkmen S, Gillessen-Kaesbach G, Meinecke P, Albrecht B, Neumann LM, Hesse V, et al. Mutations in NSD1 are responsible for Sotos syndrome, but are not a frequent finding in other overgrowth phenotypes. *Eur J Hum Genet* 2003;11:858–865. DOI:10.1038/sj.ejhg.5201050
6. Jezela-Stanek A, Kucharczyk M, Falana K, Jurkiewicz D, Mlynek M, Wicher D, et al. Malan syndrome (Sotos syndrome 2) in two patients with 19p13.2 deletion encompassing NFIX gene and novel NFIX sequence variant. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2016;160:161–167. DOI:10.5507/bp.2016.006
7. Almuriekh M, Shintani T, Fahiminiya S, Fujikawa A, Kuboyama K, Takeuchi Y, et al. Loss-of-Function Mutation in APC2 Causes Sotos Syndrome Features. *Cell Rep* 2015;10:1585–1598. DOI:10.1016/j.celrep.2015.02.011.
8. Tatton-Brown K, Douglas J, Coleman K, Baujat G, Cole TR, Das S, et al. Genotype-phenotype associations in Sotos syndrome: an analysis of 266 individuals with NSD1 aberrations. *Am J Hum Genet* 2005;77:193–204. DOI:10.1086/432082
9. Sotos JF. Sotos Syndrome 1 and 2. *Pediatr Endocrinol Rev* 2014;12:2–16
10. O’Neil Danis D, Bodamer O, Levi JR. The otolaryngologic manifestations of Sotos Syndrome 1: A systematic review. *Int J Pediatr Otorhinolaryngol* 2021;143:10649. DOI: 10.1016/j.ijporl.2021.110649
11. Adhami EJ, Cancio-Babu C. Anaesthesia in a child with Sotos syndrome. Case report. *Paediatr Anaesth* 2003;13:835–840
12. Chung JY; Kim G. Anesthetic considerations in a child with Sotos Syndrome: A case report. *Anesth Pain Med* 2017;12:240–242
13. Lane C, Milne E, Freeth M. Cognition and Behaviour in Sotos Syndrome: A Systematic Review. *PLoS One* 2016;11:e0149189. DOI:10.1371/journal.pone.0149189
14. Cole T. Congenital urological anomalies in Sotos syndrome. *Br J Urol* 1996;78:156
15. Chierichini A, Messina A. Regional anesthesia in a child with Sotos Syndrome. *Int J Immunopathol Pharmacol* 2011;24;1(s2):21–23
16. Joo-Young Chung, Gahyun Kim, Ju-Hun Park. Anesthetic considerations in a child with Sotos syndrome - A case report. *Anesth Pain Med* 2017;12: 240–242. DOI:10.17085/apm.2017.12.3.240
17. Fortin O, Vincelette C, Khan AQ. Seizures in Sotos syndrome: Phenotyping in 49 patients. *Epilepsia Open* 2021; 6:425–430. Published online 2021 Apr 9. DOI: 10.1002/epi4.12484
18. Suresh D. Posterior spinal fusion in Sotos' syndrome. *Br J Anaesth* 1991;66:728–732. DOI:10.1093/bja/66.6.728
19. Varvinski A, McGill FJ, Judd V, Hodzovic I. Sotos' syndrome ... a rare challenge? *Anaesthesia* 2001;56:809. DOI:10.1046/j.1365-2044.2001.02181-17.x.

Date last modified: **December 2021**

This recommendation was prepared by:

Author

Kim Caulfield, Consultant Anaesthesiologist, University Hospital Waterford, Ireland
kimcaulfield86@gmail.com

Patrick Wiseman, Anaesthetic Trainee, Beaumont Hospital
patrickwiseman@hotmail.com

Disclosure These authors have no financial or other competing interest to disclose. This recommendation was unfunded.

This recommendation was reviewed by:

Reviewers

Arash Afshari, Anaesthesiologist, Department of Pediatric and Obstetric Anesthesia,
University of Copenhagen, Rigshospitalet, Denmark
Arash.Afshari@regionh.dk

Cristina del Rio Peña, Pediatric Anesthesiology, 12 de Octubre Hospital, Madrid. Miembro
del Grupo de Trabajo "Enfermedades Raras y Anestesia" de la Sección Pediátrica de la
SEDAR
cristina.rio@salud.madrid.org

Disclosure The reviewers have no financial or other competing interest to disclose.

Please note that this recommendation has not been reviewed by an anaesthesiologist and a disease expert but by two anaesthesiologists instead.
